



WEBER CITY  
PROFESSIONAL CORNER  
SCOTT DANLEY D.D.S.  
STEVEN R. BROWN D.D.S. M.S.D.  
DENTWOODVILLE  
FURNER MANAGEMENT CORP.












Figure 1. Schematic representation of the experimental design. The subjects were divided into two groups: the control group and the experimental group. The control group was divided into two subgroups: the control group and the control group. The experimental group was divided into two subgroups: the experimental group and the experimental group. The control group was divided into two subgroups: the control group and the control group. The experimental group was divided into two subgroups: the experimental group and the experimental group.

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D).